arXiv:math/0608649v1 [math.NT] 26 Aug 2006

# A NOTE ON $q$-EULER NUMBERS AND POLYNOMIALS

Taekyun Kim

Abstract. The purpose of this paper is to construct $q$-Euler numbers and polynomials by using $p$-adic $q$-integral equations on $\mathbb{Z}_p$. Finally, we will give some interesting formula related to these $q$-Euler numbers and polynomials.

## §1. Introduction

The usual Bernoulli numbers are defined by

$$\sum_{k=0}^{\infty} B_k \frac{t^k}{k!} = \frac{t}{e^t-1},$$

which can be written symbolically as $e^{Bt} = \frac{t}{e^t-1}$, interpreted to mean $B^k$ must be replaced by $B_k$ when we expand on the left. This relation can also be written $e^{(B+1)t} - e^{Bt} = t$, or, if we equate power of $t$,

$$B_0 = 1, \quad (B+1)^k - B_k = \begin{cases} 1 & \text{if } k = 1 \\ 0 & \text{if } \ k > 1, \end{cases}$$

where again we must first expand and then replace $B^i$ by $B_i$, cf. [6,8,9,10,11].

Carlitz's $q$-Bernoulli numbers $\beta_k$ can be determined inductively by

$$\beta_0 = 1, \quad q(q\beta+1)^k - \beta_k = \begin{cases} 1 & \text{if } k = 1 \\ 0 & \text{if } \ k > 1, \end{cases} \tag{1}$$

with the usual convention about replacing $\beta^i$ by $\beta_{i,q}$(see [1,2,3,4,5,12]).

Carlitz also defined $q$-Euler numbers and polynomials as

$$H_0(u;q) = 1, \quad (qH+1)^k - uH_k(u;q) = 0 \quad \text{for } k \geq 1, \tag{2}$$

where $u$ is a complex number with $|u| > 1$; and for $k \geq 0$,

$$H_k(u,x;q) = (q^x H + [x]_q)^k, \quad \text{cf. } [2,3,4,9,10,11,12],$$

Typeset by $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-TEX

where $[x]_q$ is defined by $[x]_q = \frac{1-q^x}{1-q}$.

It was known that the ordinary Euler polynomials are defined by

$$\frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}, \quad (|t| < \pi), \tag{3}$$

we note that $E_n = E_n(0)$ are called $n$-th Euler numbers, cf. [6,7,8].

Let $p$ be a fixed odd prime, and let $\mathbb{C}_p$ denote the $p$-adic completion of the algebraic closure of $\mathbb{C}_p$. For $d$ a fixed positive integer $(p,d) = 1$, let

$$\begin{aligned} &X = X_d = \varprojlim_N \mathbb{Z}/dp^N\mathbb{Z}, \\ &X_1 = \mathbb{Z}_p, \\ &X^* = \underset{\substack{0<a<dp \\ (a,p)=1}}{\cup} (a + dp\mathbb{Z}_p), \\ &a + dp^N\mathbb{Z}_p = \{x \in X | x \equiv a \pmod{dp^n}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \le a < dp^N$.

The $p$-adic absolute value in $\mathbb{C}_p$ is normalized so that $|p|_p = \frac{1}{p}$. Let $q$ be variously considered as an indeterminate a complex number $q \in \mathbb{C}$, or a $p$-adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, we always assume $|q| < 1$. If $q \in \mathbb{C}_p$, we always assume $|q-1|_p < p^{-1/p-1}$, so that $q^x = \exp(x \log q)$ for $|x|_p \le 1$.

Throughout this paper we use the notation:

$$[x]_q = \frac{q^x - 1}{q - 1} = 1 + q + q^2 + \cdots + q^{x-1}.$$

We say that $f$ is uniformly differentiable function at a point $a \in \mathbb{Z}_p$ and denote this property by $f \in UD(\mathbb{Z}_p)$, if the difference quotients

$$F_f(x,y) = \frac{f(x) - f(y)}{x - y}$$

have a limit $l = f'(a)$ as $(x,y) \to (a,a)$.

For $f \in UD(\mathbb{Z}_p)$, let us start with expression

$$\frac{1}{[p^N]_q} \sum_{0 \le j < p^N} q^j f(j) = \sum_{0 \le j < p^N} f(j)\mu_q(j + p^N\mathbb{Z}_p), \tag{4}$$

representing $q$-analogue of Riemann sums for $f$, cf. [5]. The integral of $f$ on $\mathbb{Z}_p$ will be defined as limit $(n \to \infty)$ of those sums, when it exists. The $q$-Volkenborn integral of function $f \in UD(\mathbb{Z}_p)$ is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{0 \le x < p^N} f(x) q^x, \quad \text{(see [5] ).} \tag{5}$$

From the definition of $[x]_q$, we derive

$$[x]_{-q} = \frac{1-(-q)^x}{1+q} = 1 - q + q^2 - q^3 + \cdots + (-1)^{x-1}q^{x-1}.$$

In [4,5], it was known that

$$\int_{\mathbb{Z}_p} [x+y]_q^m d\mu_q(y) = \beta_{m,q}(x), \tag{6}$$

where $\beta_{m,q}(x)$ are called Carlitz's $q$-Bernoulli polynomials. By (5), we easily see that

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \frac{[2]_q}{2} \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x q^x. \tag{7}$$

The purpose of this note is to construct $q$-Euler numbers which can be uniquely determined by

$$q(qE_q+1)^n + E_{n,q} = \begin{cases} [2]_q & \text{if } n = 0 \\ 0 & \text{if } \ n > 0, \end{cases} \tag{8}$$

with the usual convention about replacing $E_q^i$ by $E_{i,q}$. From these numbers, we will derive some interesting formulae.

## §2. A note on $q$-Euler numbers and polynomials

From (7), we derive formulae as follows:

$$qI_{-q}(f_1) + I_{-q}(f) = [2]_q f(0), \tag{9}$$

where $f_1(x)$ is translation with $f_1(x) = f(x+1)$.

Let $f(x) = e^{[x]_q t}$. Then we see that

$$q\int_{\mathbb{Z}_p} e^{[x+1]_q t} d\mu_{-q}(x) + \int_{\mathbb{Z}_p} e^{[x]_q t} d\mu_{-q}(x) = [2]_q. \tag{10}$$

First, we consider the following integral :

$$\begin{aligned}\int_{\mathbb{Z}_p} e^{[x]_q t} d\mu_{-q}(x) &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\frac{1}{1-q}\right)^n \sum_{l=0}^{n} \binom{n}{l} (-1)^l \int_{\mathbb{Z}_p} q^{xl} d\mu_{-q}(x) \frac{t^n}{n!} \\ &= [2]_q \sum_{n=0}^{\infty} \left(\frac{1}{1-q}\right)^n \sum_{l=0}^{n} \binom{n}{l} (-1)^l \sum_{m=0}^{\infty} (-1)^m q^{(l+1)m} \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m e^{[m]_q t} = \sum_{n=0}^{\infty} E_{n,q} \frac{t^n}{n!}. \end{aligned} \tag{11}$$

By (11), we easily see that

$$\int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) = E_{n,q}. \tag{12}$$

Note that

$$\lim_{q \to 1} E_{n,q}(x) = E_n = \int_{\mathbb{Z}_p} x^n d\mu_{-q}(x), \quad \text{see[7]}.$$

By the same method, we note that

$$\begin{aligned}\int_{\mathbb{Z}_p} e^{[x+y]_q t} d\mu_{-q}(y) &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} [x+y]_q^n d\mu_{-q}(y) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\frac{1}{1-q}\right)^n \left( \sum_{l=0}^{n} \binom{n}{l} (-1)^l q^{lx} \int_{\mathbb{Z}_p} q^{yl} d\mu_{-q}(y) \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{n=0}^{\infty} \left(\frac{1}{1-q}\right)^n \left( \sum_{l=0}^{n} \binom{n}{l} \frac{(-1)^l q^{lx}}{1+q^{l+1}} \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} [m+x]_q^n \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m e^{[x+m]_q t} \\ &= \sum_{m=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}. \end{aligned}$$

Thus, we have

$$\int_{\mathbb{Z}_p} [x+y]_q^n d\mu_{-q}(y) = E_{n,q}(x), \quad n \geq 0. \tag{13}$$

Note that

$$\lim_{q\to 1} E_{n,q}(x) = E_n(x) = \int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(y).$$

Since

$$[x+1]_q = \frac{1-q^{x+1}}{1-q} = \frac{1-q}{1-q} + \frac{1-q^x}{1-q}q = 1 + q[x]_q. \tag{14}$$

From (10), (12) and (14), we derive

$$\begin{aligned}
[2]_q &= \sum_{n=0}^{\infty} \left( q\int_{\mathbb{Z}_p} [x+1]_q^n d\mu_{-q}(x) + \int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left( q\sum_{l=0}^{n} \binom{n}{l} q^l \int_{\mathbb{Z}_p} [x]_q^l d\mu_{-q}(x) + \int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left( q\sum_{l=0}^{n} \binom{n}{l} q^l E_{l,q} + E_{n,q} \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left( q(qE_q+1)^n + E_{n,q} \right) \frac{t^n}{n!},
\end{aligned} \tag{15}$$

with the usual convention about replacing $E_q^i$ by $E_{i,q}$.

By (15), we easily see that

$$q(qE_q+1)^n + E_{n,q} = \begin{cases} [2]_q & \text{if } n=0 \\ 0 & \text{if } n>0, \end{cases} \tag{16}$$

with the usual convention about replacing $E_q^i$ by $E_{i,q}$. When we compare Eq.(16) and Eq.(1), the Eq.(16) seems to be interesting formula. In particular, these numbers seem to be new, which are different than Carlitz's $q$-Euler numbers. From (7), we derive

$$q^n I_{-q}(f_n) + (-1)^{n-1} I_{-q}(f) = [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l), \tag{16-1}$$

where $n \in \mathbb{N}$, $f_n(x) = f(x+n)$. When $n$ is an odd positive integer, we note that

$$q^n I_{-q}(f_n) + I_{-q}(f) = [2]_q \sum_{l=0}^{n-1} (-1)^l q^l f(l). \tag{17}$$

From (17), we derive

$$q^n \int_{\mathbb{Z}_p} [x+n]_q^m d\mu_{-q}(x) + \int_{\mathbb{Z}_p} [x]_q^m d\mu_{-q}(x) = [2]_q \sum_{l=0}^{n-1} (-1)^l q^l [l]_q^m, \tag{18}$$

where $n$ is an odd positive integer. By (12) and (13), we easily see that

$$[2]_q \sum_{l=0}^{n-1} (-1)^l q^l [l]_q^m = q^n E_{m,q}(n) + E_{m,q}, \tag{19}$$

where $n$ is an odd positive integer. If $n$ is an even integer, then we have in Eq.(16-1) that

$$q^n \int_{\mathbb{Z}_p} [x+n]_q^m d\mu_{-q}(x) - \int_{\mathbb{Z}_p} [x]_q^m d\mu_{-q}(x) = [2]_q \sum_{l=0}^{n-1} (-1)^{l-1} q^l [l]_q^m. \tag{20}$$

From (12), (13) and (20), we derive

$$q^n E_{m,q}(n) - E_{m,q} = [2]_q \sum_{l=0}^{n-1} (-1)^{l-1} q^l [l]_q^m, \tag{21}$$

where $n$ is a positive even integer. It seems to be interesting to compare (19) and (21).

## References

[1] L. C. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987–1000.

[2] L. C. Carlitz, *q-Bernoulli and Eulerian numbers*, Trans. Amer. Math. Soc. **76** (1954), 332–350.

[3] M. Cenkci, M. Can and V. Kurt, *p-adic interpolation functions and Kummer-type congruences for q-twisted Euler numbers*, Advan. Stud. Contemp. Math. **9** (2004), 203–216.

[4] M. Cenkci, M. Can, *Some results on q-analogue of the Lerch zeta function*, Adv. Stud. Contemp. Math. **12** (2006), 213–223.

[5] T. Kim, *q− Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.

[6] T. Kim, *A note on p-adic invariant integral in the rings of p-adic integers*, Advan. Stud. Contemp. Math. **13** (2006), 95–99.

[7] T. Kim, *q-generalized Euler numbers and polynomials*, Russian J. Math. Phys. **13** (2006), 293-298.

[8] T. Kim, *A note on q-Euler and Genocchi numbers*, Proc. Japan Acad. Ser. A **77** (2001), 139-141.

[9] N. Koblitz, *q-Bernoulli numbers*, J. Number Theory **14** (1982), 332–339.

[10] Y. Simsek, *Theorems on twisted L-function and twisted Bernoulli numbers*, Advan. Stud. Contem. Math. **11** (2005), 205–218.

[11] Y. Simsek, *q-Dedekind type sums related to q-zeta function and basic L-series*, J. Math. Anal. Appl. **318** (2006), 333–351.

[12] H. M. Srivastava, T. Kim and Y. Simsek, *q-Bernoulli numbers and polynomials associated with multiple q-zeta functions and basic L-series*, Russ. J. Math. Phys. **12** (2005), 241–268.

Taekyun Kim
EECS, Kyungpook National University, Taegu 702-701, S. Korea
e-mail: tkim@knu.ac.kr; tkim64@hanmail.net